



WHAT WE KNOW

Psychosocial Treatment for Children and Adolescents with AD/HD*

P psychosocial treatment is a critical part of treatment for attention-deficit/hyperactivity disorder

(AD/HD) in children and adolescents. The scientific literature, the National Institute of Mental Health, and many professional organizations agree that behaviorally oriented psychosocial treatments—also called behavior

therapy or behavior modification—and stimulant medication have a solid base of scientific evidence demonstrating their effectiveness. Behavior modification is the only nonmedical treatment for AD/HD with a large scientific evidence base.

Treating AD/HD in children often involves medical, educational and behavioral interventions. This comprehensive approach to treatment is called “multimodal” and consists of parent and child education about diagnosis and treatment, behavior management techniques, medication, and school programming and supports. The severity and type of AD/HD may be factors in deciding which components are necessary. Treatment should be tailored to the unique needs of each child and family.

* This fact sheet should be read together with What We Know #3, “Managing Medication for Children and Adolescents with AD/HD.”

This fact sheet will:

- define behavior modification
- describe effective parent training, school interventions and child interventions
- discuss the relationship between behavior modification and stimulant medication in treating children and adolescents with AD/HD

WHY USE PSYCHOSOCIAL TREATMENTS?

Behavioral treatment for AD/HD is important for several reasons. First, children with AD/HD face problems in daily life that go well beyond their symptoms of inattentiveness, hyperactivity and impulsivity, including poor academic performance and behavior at school, poor relationships with peers and siblings, failure to obey adult requests, and poor relationships with their parents. These problems are extremely important because they predict how children with AD/HD will do in the long run.

How a child with AD/HD will do in adulthood is best predicted by three things—(1) whether his or her parents use effective parenting skills, (2) how he or she gets along with other children, and (3) his or her success in school¹. Psychosocial treatments are effective in treating these important domains. Second, behavioral treatments teach skills to parents and teachers that help them deal with children with AD/HD. They also teach skills to children with AD/HD that will help them overcome their impairments. Learning these skills is especially important because AD/HD is a chronic condition and these skills will be useful throughout the children's lives².

Behavioral treatments for AD/HD should be started as soon as the child receives a diagnosis. There are behavioral interventions that work well for preschoolers, elementary-age students, and teenagers with AD/HD, and there is consensus that starting early is better than starting later. Parents, schools, and practitioners should not put off beginning effective behavioral treatments for children with AD/HD^{3,4}.

WHAT IS BEHAVIOR MODIFICATION?

With behavior modification, parents, teachers and children learn specific techniques and skills from a therapist, or an educator experienced in the approach, that will help improve children's behavior. Parents and teachers then use the skills in their daily interactions with their children with AD/HD, resulting in

improvement in the children's functioning in the key areas noted above. In addition, the children with AD/HD use the skills they learn in their interactions with other children.

Behavior modification is often put in terms of ABCs: Antecedents (things that set off or happen before behaviors), Behaviors (things the child does that parents and teachers want to change), and Consequences (things that happen after behaviors). In behavioral programs, adults learn to change antecedents (for example, how they give commands to children) and consequences (for example, how they react when a child obeys or disobeys a command) in order to change the child's

“Parent, teacher and child interventions should be carried out at the same time to get the best result...”

behavior (that is, the child's response to the command). By consistently changing the ways that they respond to children's behaviors, adults teach the children new ways of behaving.

Parent, teacher and child interventions should be carried out at the same time to get the best results^{5,6}. The following four points should be incorporated into all three components of behavior modification:

1. Start with goals that the child can achieve in small steps.
2. Be consistent—across different times of the day, different settings, and different people.
3. Implement behavioral interventions over the long haul—not just for a few months.
4. Teaching and learning new skills take time, and children's improvement will be gradual.

Parents who want to try a behavioral approach with their children should learn what distinguishes behavior modification from other approaches so they can recognize effective behavioral treatment and be confident that what the therapist is offering will improve their child's functioning. Many psychotherapeutic treatments have not been proven to work for children with AD/HD. Traditional individual therapy, in which a child spends time with a therapist or school counselor talking about

his or her problems or playing with dolls or toys, is not behavior modification. Such “talk” or “play” therapies do not teach skills and have not been shown to work for children with AD/HD^{7,8}.

HOW DOES A BEHAVIOR MODIFICATION PROGRAM BEGIN?

The first step is identifying a mental health professional who can provide behavioral therapy. Finding the right professional may be difficult for some families, especially for those that are economically disadvantaged or socially or geographically isolated. Families should ask their primary care physicians for a referral or contact their insurance company for a list of providers who participate in the insurance plan, though health insurance may not cover the costs of the kind of intensive treatment that is most helpful. Other sources of referrals include professional associations and hospital and university AD/HD centers (visit www.help4adhd.org for a list).

The mental health professional begins with a complete evaluation of the child’s problems in daily life, including home, school (both behavioral and academic), and social settings. Most of this information comes from parents and teachers. The therapist also meets with the child to get a sense of what the child is like. The evaluation should result in a list of target areas for treatment. Target areas—often called target behaviors—are behaviors in which change is desired, and if changed, will help improve the child’s functioning/impairment and long-term outcome.

Target behaviors can be either negative behaviors that need to stop or new skills that need to be developed. That means that the areas targeted for treatment will typically not be the symptoms of AD/HD—overactivity, inattention and impulsivity—but rather the specific problems that those symptoms may cause in daily life. Common classroom target behaviors include “completes assigned work with 80 percent accuracy” and “follows classroom rules.” At home, “plays well with siblings (that is, no fights)” and “obeys parent requests or commands” are common target behaviors. (Lists of common target behaviors in school, home and peer settings can be downloaded in Daily Report Card packets at <http://wings.buffalo.edu/adhd>.)

After target behaviors are identified, similar behavioral interventions are implemented at home and at school. Parents and teachers learn and establish programs in which the environmental antecedents (the A’s) and

consequences (the C’s) are modified to change the child’s target behaviors (the B’s). Treatment response is constantly monitored, through observation and measurement, and the interventions are modified when they fail to be helpful or are no longer needed.

PARENT TRAINING

Behavioral parent training programs have been used for many years and have been found to be very effective⁹⁻¹⁹.

Although many of the ideas and techniques taught in behavioral parent training are common sense parenting techniques, most parents need careful teaching and support to learn parenting skills and use them consistently. It is very difficult for parents to buy a book, learn behavior modification, and implement an effective program on their own. Help from a professional is often necessary. The topics covered in a typical series of parent training sessions include the following:

- Establishing house rules and structure
- Learning to praise appropriate behaviors (praising good behavior at least five times as often as bad behavior is criticized) and ignoring mild inappropriate behaviors (choosing your battles)
- Using appropriate commands
- Using “when...then” contingencies (withdrawing rewards or privileges in response to inappropriate behavior)
- Planning ahead and working with children in public places
- Time out from positive reinforcement (using time outs as a consequence for inappropriate behavior)
- Daily charts and point/token systems with rewards and consequences
- School-home note system for rewarding behavior at school and tracking homework^{20,21}

Some families can learn these skills quickly in the course of 8-10 meetings, while other families—often those with the most severely affected children—require more time and energy.

Parenting sessions usually involve an instructional book or videotape on how to use behavioral management procedures with children. The first session is often devoted to an overview of the diagnosis, causes, nature, and prognosis of AD/HD. Next, parents learn a variety of techniques, which they may already be using at home but not as consistently or correctly as needed. Parents then go home and implement what they have learned

in sessions during the week, and return to the parenting session the following week to discuss progress, solve problems, and learn a new technique.

Parent training can be conducted in groups or with individual families. Individual sessions often are implemented when a group is not available or when the family would benefit from a tailored approach that includes the child in sessions. This kind of treatment is called behavioral family therapy. The number of family therapy sessions varies depending on the severity of the problems^{22,23,24}. CHADD offers a unique educational program to help parents and individuals navigate the challenges of AD/HD across the lifespan. Information about CHADD's "Parent to Parent" program can be found by visiting CHADD's Web site at www.chadd.org. Click on "Conferences and Training" and then "Parent to Parent Program."

When the child involved is a teenager, parent training is slightly different. Parents are taught behavioral techniques that are modified to be age-appropriate for adolescents. For example, time out is a consequence that is not effective with teenagers; instead, loss of privileges (such as having the car keys taken away) or assignment of work chores would be more appropriate. After parents

"Interventions for peer relationships (how the child gets along with other children) are a critical component of treatment for children with AD/HD."

have learned these techniques, the parents and teenager typically meet with the therapist together to learn how to come up with solutions to problems on which they all agree. Parents negotiate for improvements in the teenagers' target behaviors (such as better grades in school) in exchange for rewards that they can control (such as allowing the teenager to go out with friends). The give and take between parents and teenager in these sessions is necessary to motivate the teenager to work with the parents in making changes in his or her behavior.

Applying these skills with children and teens with AD/HD takes a lot of hard work on the part of parents.

However, the hard work pays off. Parents who master and consistently apply these skills will be rewarded with a child who behaves better and has a better relationship with parents and siblings.

SCHOOL INTERVENTIONS

As is the case with parent training, the techniques used to manage AD/HD in the classroom have been used for some time and are considered effective^{25,26,27,28,29,30,31}. Many teachers who have had training in classroom management are quite expert in developing and implementing programs for students with AD/HD. However, because the majority of children with AD/HD are not enrolled in special education services, their teachers will most often be regular education teachers who may know little about AD/HD or behavior modification and will need assistance in learning and implementing the necessary programs. There are many widely available handbooks, texts and training programs that teach classroom behavior management skills to teachers. Most of these programs are designed for regular or special education classroom teachers who also receive training and guidance from school support staff or outside consultants. Parents of children with AD/HD should work closely with the teacher to support efforts in implementing classroom programs. (To read more about typical classroom behavioral management procedures, please see Appendix A.)

Managing teenagers with AD/HD in school is different from managing children with AD/HD. Teenagers need to be more involved in goal planning and implementation of interventions than do children. For example, teachers expect teenagers to be more responsible for belongings and assignments. They may expect students to write assignments in weekly planners rather than receive a daily report card. Organizational strategies and study skills therefore need to be taught to the adolescent with AD/HD. Parent involvement with the school, however, is as important at the middle and high school levels as it is in elementary school. Parents will often work with guidance counselors rather than individual teachers, so that the guidance counselor can coordinate intervention among the teachers.

CHILD INTERVENTIONS

Interventions for peer relationships (how the child gets along with other children) are a critical component

of treatment for children with AD/HD. Very often, children with AD/HD have serious problems in peer relationships^{32,33,34,35}. Children who overcome these problems do better in the long run than those who continue to have problems with peers³⁶. There is scientific basis for child-based treatments for AD/HD that focus on peer relationships. These treatments usually occur in group settings outside of the therapist's office.

There are five effective forms of intervention for peer relationships:

1. systematic teaching of social skills³⁷
2. social problem solving^{38,39,40}
3. teaching other behavioral skills often considered important by children, such as sports skills and board game rules⁴¹
4. decreasing undesirable and antisocial behaviors^{42,43}
5. developing a close friendship

There are several settings for providing these interventions to children, including groups in office clinics, classrooms, small groups at school, and summer camps. All of the programs use methods that include coaching, use of examples, modeling, role-playing, feedback, rewards and consequences, and practice. It is best if these child-directed treatments are used when a parent is participating in parent training and school personnel are conducting an appropriate school intervention^{44,45,46,47}. When parent and school interventions are integrated with child-focused treatments, problems getting along with other children (such as being bossy, not taking turns, and not sharing) that are being targeted in the child treatments are also included as target behaviors in the home and school programs so that the same behaviors are being monitored, prompted and rewarded in all three settings.

Social skills training groups are the most common form of treatment, and they typically focus on the systematic teaching of social skills. They are typically conducted at a clinic or in school in a counselor's office for 1-2 hours on a weekly basis for 6-12 weeks. Social skills groups with children with AD/HD are only effective when they are used with parent and school interventions and rewards and consequences to reduce disruptive and negative behaviors^{48,49,50,51,52}.

There are several models for working on peer relationships in the school setting that integrate several of the interventions listed above. They combine skills training with a major focus on decreasing negative and disruptive behavior and are typically conducted

by school staff. Some of these programs are used with individual children (for instance, token programs in the classroom or at recess)^{53,54} and some are schoolwide (such as peer mediation programs)^{55,56}.

Generally, the most effective treatments involve helping children get along better with other children. Programs in which children with AD/HD can work on peer problems in classroom or recreational settings are the most effective^{57,58}. One model involves establishing a summer camp for children with AD/HD in which child-based management of peer problems and academic difficulties are integrated with parent training^{59,60,61}. All five forms of peer intervention are incorporated in a 6-8 week program that runs for 6-9 hours on weekdays. Treatment is conducted in groups, with recreational activities (e.g., baseball, soccer) for the majority of the day, along with two hours of academics. One major focus is teaching skills in and knowledge of sports to the children. This is combined with intensive practice in social and problem-solving skills, good team work, decreasing negative behaviors, and developing close friendships.

Some approaches to child-based treatment for peer problems fall somewhere between clinic-based programs and intensive summer camps. Versions of both are conducted on Saturdays during the school year or after school. These involve 2-3 hour sessions in which children engage in recreational activities that integrate many of the forms of social skills intervention.

Finally, preliminary research suggests that having a best friend may have a protective effect on children with difficulties in peer relations as they develop through childhood and into adolescence^{62,63}. Researchers have developed programs that help children with AD/HD build at least one close friendship. These programs always begin with the other forms of intervention described above and then add having the families schedule monitored play dates and other activities for their child and another child with whom they are attempting to foster a friendship.

It is important to emphasize that simply inserting a child with AD/HD in a setting where there is interaction with other children—such as Scouts, Little League or other sports, day care, or playing in the neighborhood without supervision—is not effective treatment for peer problems. Treatment for peer problems is quite complex and involves combining careful instruction in social and problem-solving skills with supervised practice in peer settings in which children receive rewards and

consequences for appropriate peer interactions. It is very difficult to intervene in the peer domain, and Scout leaders, Little League coaches, and day-care personnel are typically not trained to implement effective peer interventions.

WHAT ABOUT COMBINING PSYCHOSOCIAL APPROACHES WITH MEDICATION?

Numerous studies over the last 30 years show that both medication and behavioral treatment are effective in improving AD/HD symptoms. Short-term treatment studies that compared medication to behavioral treatment have found that medication alone is more effective in treating AD/HD symptoms than behavioral treatment alone. In some cases, combining the two approaches resulted in slightly better results.

The best-designed long-term treatment study—the Multimodal Treatment Study of Children with AD/HD (MTA)—was conducted by the National Institute of Mental Health. The MTA studied 579 children with AD/HD-combined type over a 14-month period. Each child received one of four possible treatments: medication management, behavioral treatment, a combination of the two, or the usual community care. The results of this landmark study were that children who were treated with medication alone, which was carefully managed and individually tailored, and children who received both medication and behavioral treatment experienced the greatest improvements in their AD/HD symptoms^{64,55}.

Combination treatment provided the best results in improving AD/HD and oppositional symptoms and in other areas of functioning, such as parenting and academic outcomes⁶⁶. Overall, those who received closely monitored medication management had greater improvement in their AD/HD symptoms than children who received either intensive behavioral treatment without medication or community care with less carefully monitored medication. It is unclear whether children with the inattentive type will show the same pattern of response to behavioral interventions and medication as have children with combined type.

Some families may choose to try stimulant medication first, while others may be more comfortable beginning with behavioral therapy. Another option is to incorporate both approaches into the initial treatment plan. The combination of the two modalities may enable

the intensity (and expense) of behavioral treatments and the dose of medication to be reduced^{67,68,69,70}.

A growing number of physicians believe that stimulant medication should not be used as the only intervention and should be combined with parent training and classroom behavioral interventions^{71,72}. In the end, each family has to make treatment decisions based on the available resources and what makes the best sense for the particular child. No one treatment plan is appropriate for everyone.

WHAT IF THERE ARE OTHER PROBLEMS IN ADDITION TO AD/HD?

There are evidence-based behavioral treatments for problems that can co-exist with AD/HD, such as anxiety⁷³ and depression⁷⁴. Just as play therapy and other non-behaviorally based therapies are not effective for AD/HD, they have not been documented to be effective for the conditions that often occur with AD/HD.

SUGGESTED READING FOR PROFESSIONALS

Barkley, R.A. (1987). *Defiant children: A clinician's manual for parent training*. New York: Guilford.

Barkley, R.A., & Murphy, K.R. (1998). *Attention-deficit hyperactivity disorder: A clinical workbook*. (2nd ed.). New York: Guilford.

Chamberlain, P. & Patterson, G.R. (1995). Discipline and child compliance in parenting. In M. Bornstein (Ed.), *Handbook of parenting: Vol. 4. Applied and practical parenting*. (pp. 205–225). Mahwah, NJ: Lawrence Erlbaum Associates.

Coie, J.D., & Dodge, K.A. (1998). Aggression and antisocial behavior. In W. Damon (Series Ed.) & N. Eisenberg (Vol. Ed.), *Handbook of child psychology: Vol. 3. Social, emotional, and personality development*. (5th ed., pp.779–862). New York: John Wiley & Sons, Inc.

Dendy, C. (2000). *Teaching teens with ADD and ADHD: A quick reference guide for teachers and parents*. Bethesda, MD: Woodbine House.

DuPaul, G.J., & Stoner, G. (2003). *AD/HD in the schools: Assessment and intervention strategies* (2nd ed.). New York: Guilford.

Forehand, R., & Long, N. (2002). *Parenting and the strong-willed child*. Chicago, IL: Contemporary Books.

Hembree-Kigin, T.L., & McNeil, C.B. (1995). *Parent-child interaction therapy: A step-by-step guide for clinicians*. New York: Plenum Press.

Kazdin, A.E. (2001). *Behavior modification in applied settings*. (6th ed.). Belmont, CA: Wadsworth/Thomson Learning.

Kendall, P.C. (2000). *Cognitive-behavioral therapy for anxious children: Therapist manual* (2nd ed.). Ardmore, PA: Workbook Publishing.

Martin, G., & Pear, J. (2002). *Behavior modification: What it is and how to do it*. (7th ed.). Upper Saddle River, NJ: Prentice-Hall, Inc.

McFayden-Ketchum, S.A. & Dodge, K.A. (1998). Problems in social relationships. In E.J. Mash & R.A. Barkley (Eds.). *Treatment of childhood disorders*. (2nd ed., pp 338–365). New York: Guilford Press.

Mrug, S., Hoza, B., & Gerdes, A.C. (2001). Children with attention-deficit/hyperactivity disorder: Peer relationships and peer-oriented interventions. In D.W. Nangle & C.A. Erdley (Eds.). *The role of friendship in psychological adjustment: New directions for child and adolescent development* (pp. 51–77). San Francisco: Jossey-Bass.

Pelham, W.E., & Fabiano, G.A. (2000). Behavior modification. *Psychiatric Clinics of North America*, 9, 671–688.

Pelham, W.E., Fabiano, G.A., Gnagy, E.M., Greiner, A.R., & Hoza, B. (in press). Comprehensive psychosocial treatment for AD/HD. In E. Hibbs & P. Jensen (Eds.), *Psychosocial treatments for child and adolescent disorders: Empirically based strategies for clinical practice*. New York: APA Press.

Pelham, W.E., Greiner, A.R., & Gnagy, E.M. (1997). *Children's summer treatment program manual*. Buffalo, NY: Comprehensive Treatment for Attention Deficit Disorders.

Pelham, W. E., Wheeler, T., & Chronis, A. (1998). Empirically supported psychosocial treatments for attention deficit hyperactivity disorder. *Journal of Clinical Child Psychology*, 27, 190-205.

Pfiffner, L.J. (1996). *All about AD/HD: The complete practical guide for classroom teachers*. New York: Scholastic Professional Books.

Rief, S.F., & Heimburge, J.A. (2002). *How to reach and teach ADD/AD/HD children: Practical techniques, strategies, and interventions for helping children with attention problems and hyperactivity*. San Francisco: Jossey-Bass.

Robin, A.L. (1998). *AD/HD in adolescents: Diagnosis and treatment*. New York: Guilford Press.

Walker, H.M., Colvin, G., & Ramsey, E. (1995). *Antisocial behavior in school: Strategies and best practices*. Pacific Grove, CA: Brooks/Cole Publishing Company.

Walker, H.M., & Walker, J.E. (1991). *Coping with noncompliance in the classroom: A positive approach for teachers*. Austin, TX: ProEd.

Wielkiewicz, R.M. (1995). *Behavior management in the schools: Principles and procedures* (2nd ed.). Boston: Allyn and Bacon.

SUGGESTED READING FOR PARENTS/CAREGIVERS

Barkley, R.A. (1987). *Defiant children: Parent-teacher assignments*. New York: Guilford Press.

Barkley, R.A. (1995). *Taking charge of AD/HD: The complete, authoritative guide for parents*. New York: Guilford.

Dendy, C. (1995). *Teenagers with ADD: A parents' guide*. Bethesda, MD: Woodbine House

Forehand, R. & Long, N. (2002) *Parenting and the strong-willed child*. Chicago, IL: Contemporary Books.

Greene, R. (2001). *The explosive child: A new approach for understanding and parenting easily frustrated, chronically inflexible children*. New York: Harper Collins.

Forgatch, M., & Patterson, G. R. (1989). *Parents and adolescents living together: Part 2: Family problem solving*. Eugene, OR: Castalia.

Kelley, M. L. (1990). *School-home notes: Promoting children's classroom success*. New York: Guilford Press.

Patterson, G.R., & Forgatch, M. (1987). *Parents and adolescents living together: Part 1: The basics*. Eugene, OR: Castalia.

Phelan, T. (1991). *Surviving your adolescents*. Glen Ellyn, IL: Child Management.

INTERNET RESOURCES

Center for Children and Families, University at Buffalo, <http://wings.buffalo.edu/adhd>

Comprehensive Treatment for Attention Deficit Disorder, <http://ctadd.net>

MODEL PROGRAMS

The Incredible Years

www.incredibleyears.com

Triple P: Positive Parenting Program

www.triplep.net

The Early Risers Program

August, G.J., Realmuto, G.M., Hektner, J.M., & Bloomquist, M.L. (2001). An integrated components preventive intervention for aggressive elementary school children: The Early Risers Program. *Journal of Consulting and Clinical Psychology*, 69, 614–626.

CLASS (Contingencies for Learning Academic and Social Skills)

Hops, H., & Walker, H.M. (1988). *CLASS: Contingencies for Learning Academic and Social Skills manual*. Seattle, WA: Educational Achievement Systems.

RECESS (Reprogramming Environmental Contingencies for Effective Social Skills)

Walker, H.M., Hops, H., & Greenwood, C.R. (1992). *RECESS manual*. Seattle, WA; Educational Achievement Systems.

Peabody Classwide Peer Tutoring Reading Methods

Mathes, P. G., Fuchs, D., Fuchs, L.S., Henley, A.M., & Sanders, A. (1994). Increasing strategic reading practice with Peabody Classwide Peer Tutoring. *Learning Disabilities Research and Practice*, 9, 44–48.

Mathes, P.G., Fuchs, D., & Fuchs, L.S. (1995). Accommodating diversity through Peabody Classwide Peer Tutoring. *Intervention in School and Clinic, 31*, 46–50.

COPE (Community Parent Education Program)

Cunningham, C. E., Cunningham, L. J., & Martorelli, V. (1997). *Coping with conflict at school: The collaborative student mediation project manual*. Hamilton, Ontario: COPE Works.

REFERENCES

- Hinshaw, S. (2002). Is ADHD an Impairing Condition in Childhood and Adolescence?. In P.S. Jensen & J.R. Cooper (Eds.), *Attention deficit hyperactivity disorder: State of the science, best practices* (pp. 5-1–5-21). Kingston, N.J.: Civic Research Institute.
- Pelham, W.E., Wheeler, T., & Chronis, A. (1998). Empirically supported psychosocial treatments for attention deficit hyperactivity disorder. *Journal of Clinical Child Psychology, 27*, 190–205.
- Webster-Stratton, C., Reid, M.J., & Hammond, M. (2001). Social skills and problem solving training for children with early-onset conduct problems: who benefits? *Journal of Child Psychology and Psychiatry, 42*, 943–952.
- August, G.J., Realmuto, G.M., Hektner, J.M., & Bloomquist, M.L. (2001). An integrated components preventive intervention for aggressive elementary school children: The Early Risers Program. *Journal of Consulting and Clinical Psychology, 69*, 614–626.
- American Academy of Pediatrics. (2001). Clinical practice guideline: Treatment of the school-aged child with attention-deficit/hyperactivity disorder. *Pediatrics, 108*, 1033–1044.
- U.S. Department of Health and Human Services (DHHS). (1999). *Mental Health: A Report of the Surgeon General*. Washington, DC: DHHS.
- Abikoff, H. (1987). An evaluation of cognitive behavior therapy for hyperactive children. In B.B. Lahey & A.E. Kazdin (Eds.), *Advances in clinical child psychology* (pp. 171–216). New York: Plenum Press.
- Abikoff, H. (1991). Cognitive training in ADHD children: Less to it than meets the eye. *Journal of Learning Disabilities, 24*, 205–209.
- Anastopoulos, A.D., Shelton, T.L., DuPaul, G.J., & Guevremont, D.C. (1993). Parent training for attention deficit hyperactivity disorder: Its impact on child and parent functioning. *Journal of Abnormal Child Psychology, 21*, 581–596.
- Brestan, E.V., & Eyberg, S.M. (1998). Effective psychosocial treatments of conduct-disordered children and adolescents: 29 years, 82 studies, and 5272 kids. *Journal of Clinical Child Psychology, 27*, 180–189.
- Cunningham, C.E., Bremner, R.B., & Boyle, M. (1995). Large group community-based parenting programs for families of preschoolers at risk for disruptive behaviour disorders: Utilization, cost effectiveness, and outcome. *Journal of Child Psychology and Psychiatry, 36*, 1141–1159.
- Dubey, D.R., O’Leary, S., & Kaufman, K.F. (1983). Training parents of hyperactive children in child management: A comparative outcome study. *Journal of Abnormal Child Psychology, 11*, 229–246.
- Hartman, R.R., Stage, S.A., & Webster-Stratton, C. (2003). A growth curve analysis of parent training outcomes: Examining the influence of child risk factors (inattention, impulsivity, and hyperactivity problems), parental and family risk factors. *Journal of Child Psychology & Psychiatry & Allied Disciplines, 44*, 388–398.
- McMahon, R.J. (1994). Diagnosis, assessment, and treatment of externalizing problems in children: The role of longitudinal data. *Journal of Consulting and Clinical Psychology, 62*, 901–917.
- Patterson, G.R., & Forgatch, M. (1987). *Parents and adolescents living together, part 1: The basics*. Eugene, OR: Castalia.
- Pisterman, S., McGrath, P.J., Firestone, P., Goodman, J.T., Webster, I., & Mallory, R. (1989). Outcome of parent-mediated treatment of preschoolers with attention deficit disorder with hyperactivity. *Journal of Consulting and Clinical Psychology, 57*, 636–643.
- Pisterman, S., McGrath, P.J., Firestone, P., Goodman, J.T., Webster, I., & Mallory, R. (1992). The effects of parent training on parenting stress and sense of competence. *Canadian Journal of Behavioural Science, 24*, 41–58.
- Pollard, S., Ward, E.M., & Barkley, R.A. (1983). The effects of parent training and Ritalin on the parent-child interactions of hyperactive boys. *Child and Family Therapy, 5*, 51–69.
- Stubbe, D.E., & Weiss, G. Psychosocial interventions: Individual psychotherapy with the child, and family interventions. *Child and Adolescent Psychiatric Clinics of North America, 9*, 663–670.
- Kelley, M.L. (1990). *School-home notes: Promoting children’s classroom success*. New York: Guilford Press.
- Kelley, M.L., & McCain, A.P. (1995). Promoting academic performance in inattentive children: the relative efficacy of school-home notes with and without response cost. *Behavior Modification, 19*, 357–375.
- Barkley, R.A., Guevremont, D.C., Anastopoulos, A.D., & Fletcher, K.E. (1992). A comparison of three family therapy programs for treating family conflicts in adolescents with attention-deficit hyperactivity disorder. *Journal of Consulting and Clinical Psychology, 60*, 450–462.
- Everett, C.A., & Everett, S.V. (1999). *Family therapy for ADHD: Treating children, adolescents, and adults*. New York: Guilford Press.
- Northey, Jr., W.F., Wells, K.C., Silverman, W.K., & Bailey, C.E. Childhood behavioral and emotional disorders. *Journal of Marital and Family Therapy, 29*, 523–545.
- Abramowitz, A.J., & O’Leary, S.G. (1991). Behavioral interventions for the classroom: Implications for student with ADHD. *School Psychology Review, 20*, 220–234.

26. Ayllon, T., Layman, D., & Kandel, H.J. (1975). A behavioral-educational alternative to drug control of hyperactive children. *Journal of Applied Behavior Analysis*, 8, 137–146.
27. DuPaul, G.J., & Eckert, T.L. (1997). The effects of school-based interventions for attention deficit hyperactivity disorder: A meta-analysis. *School Psychology Review*, 26, 5–27.
28. Gittelman, R., Abikoff, H., Pollack, E., Klein, D. F., Katz, S., & Mattes, J. (1980). A controlled trial of behavior modification and methylphenidate in hyperactive children. In C. K. Walen & B. Henker (Eds.), *Hyperactive children: The social ecology of identification and treatment* (pp. 221–243). New York: Academic Press.
29. O’Leary, K.D., Pelham, W.E., Rosenbaum, A., & Price, G. (1976). Behavioral treatment of hyperkinetic children: An experimental evaluation of its usefulness. *Clinical Pediatrics*, 15, 510–514.
30. Pelham, W.E., Schnedler, R.W., Bender, M.E., Miller, J., Nilsson, D., Budrow, M., et al. (1988). The combination of behavior therapy and methylphenidate in the treatment of hyperactivity: A therapy outcome study. In L. Bloomingdale (Ed.), *Attention deficit disorders* (pp. 29–48). London: Pergamon.
31. Pfiffner, L.J., & O’Leary, S.G. (1993). School-based psychological treatments. In J.L. Matson (Ed.), *Handbook of hyperactivity in children* (pp. 234–255). Boston: Allyn & Bacon.
32. Bagwell, C.L., Molina, B.S., Pelham, Jr., W.E., & Hoza, B. (2001). Attention-deficit hyperactivity disorder and problems in peer relations: Predictions from childhood to adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry*, 40, 1285–1292.
33. Blachman, D.R., & Hinshaw, S.P. (2002). Patterns of friendship among girls with and without attention-deficit/hyperactivity disorder. *Journal of Abnormal Child Psychology*, 30, 625–640.
34. Hodgens, J.B., Cole, J., & Boldizar, J. (2000). Peer-based differences among boys with ADHD. *Journal of Clinical Child Psychology*, 29, 443–452.
35. McFayden-Ketchum, S.A., & Dodge, K.A. (1998). Problems in social relationships. In E.J. Mash & R.A. Barkley (Eds.), *Treatment of childhood disorders* (2nd ed., pp 338–365). New York: Guilford Press.
36. Woodward, L.J., & Fergusson, D.M. (2000). Childhood peer relationship problems and later risks of educational underachievement and unemployment. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 41, 191–201.
37. Webster-Stratton, C., Reid, J., & Hammond, M. (2001). Social skills and problem-solving training for children with early-onset conduct problems: Who benefits?. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 42, 943–52.
38. Houk, G.M., King, M.C., Tomlinson, B., Vrabell, A., & Weeks, K. (2002). Small group intervention for children with attention disorders. *Journal of School Nursing*, 18, 196–200.
39. Kazdin, A.E., Esveldt-Dawson, K., French, N.H., & Unis, A.S. (1987). Problem-solving skills training and relationship therapy in the treatment of antisocial child behavior. *Journal of Consulting and Clinical Psychology*, 55, 76–85.
40. Kazdin, A.E., Bass, D., Siegel, T., Thomas, C. (1989). Cognitive-behavioral therapy and relationship therapy in the treatment of children referred for antisocial behavior. *Journal of Consulting and Clinical Psychology*, 57, 522–535.
41. American Academy of Child and Adolescent Psychiatry. (1997). Practice parameters for the assessment and treatment of children, adolescents, and adults with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(Suppl. 10), 85–121.
42. Walker, H.M., Colvin, G., & Ramsey, E. (1995). *Antisocial behavior in school: Strategies and best practices*. Pacific Grove, CA: Brooks/Cole Publishing Company.
43. Coie, J.D., & Dodge, K.A. (1998). Aggression and antisocial behavior. In W. Damon (Series Ed.) & N. Eisenberg (Vol. Ed.), *Handbook of child psychology: Vol. 3. Social, emotional, and personality development*. (5th ed., pp.779–862). New York: John Wiley & Sons, Inc.
44. MTA Cooperative Group. (1999). A 14-Month randomized clinical trial of treatment strategies for attention-deficit/hyperactivity disorder. *Archives of General Psychiatry*, 56, 1073–1086.
45. MTA Cooperative Group. (1999). Moderators and mediators of treatment response for children with attention-deficit/hyperactivity disorder. *Archives of General Psychiatry*, 56, 1088–1096.
46. Richters, J.E., Arnold, L.E., Jensen, P.S., Abikoff, H., Conners, C.K., Greenhill, L.L., et al. (1995). NIMH collaborative multisite multimodal treatment study of children with ADHD: I. Background and rationale. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34, 987–1000.
47. Webster-Stratton, C., Reid, J., & Hammond, M. (2004). Treating children with early-onset conduct problems: Intervention outcomes for parent, child, and teacher training. *Journal of Clinical Child and Adolescent Psychology*, 33, 105–124.
48. Bierman, K L., Miller, C.L., & Stabb, S.D. (1987). Improving the social behavior and peer acceptance of rejected boys: Effects of social skill training with instructions and prohibitions. *Journal of Consulting and Clinical Psychology*, 55, 194–200.
49. Hinshaw, S.P., Henker, B., & Whalen, C.K. (1984). Self-control in hyperactive boys in anger-inducing situations: Effects of cognitive-behavioral training and methylphenidate. *Journal of Abnormal Child Psychology*, 12, 55–77.
50. Kavale, K.A., Mathur, S. R., Forness, S.R., Rutherford, R.G., & Quinn, M.M. (1997). The effectiveness of social skills training for students with emotional or behavioral disorders: A meta-analysis. In T.E. Scruggs & M.A. Mastropieri (Eds.), *Advances in learning and behavioral disabilities* (Vol. 11, pp. 1–26). Greenwich, CT: JAI.
51. Kavale, K.A., Forness, S.R., & Walker, H.M. (1999). Interventions for oppositional defiant disorder and conduct disorder in the schools. In H. Quay & A. Hogan (Eds.), *Handbook of disruptive behavior disorders* (pp. 441–454). New York: Kluwer.

52. Pfiffner, L.J., & McBurnett, K. (1997). Social skills training with parent generalization: Treatment effects for children with attention deficit disorder. *Journal of Consulting & Clinical Psychology, 65*, 749–757.
53. Pfiffner, L.J. (1996). *All about ADHD: The complete practical guide for classroom teachers*. New York: Scholastic Professional Books.
54. Abramowitz, A.J. (1994). Classroom interventions for disruptive behavior disorder. *Child and Adolescent Psychiatric Clinics of North America, 3*, 343–360.
55. Cunningham, C.E., & Cunningham, L.J. (1995). Reducing playground aggression: Student mediation programs. *ADHD Report, 3*(4), 9–11.
56. Cunningham, C.E., Cunningham, L.J., Martorelli, V., Tran, A., Young, J., & Zacharias, R. (1998). The effects of primary division, student-mediated conflict resolution programs on playground aggression. *Journal of Child Psychology and Psychiatry and Allied Disciplines, 39*, 653–662.
57. Conners, C.K., Wells, K.C., Erhardt, D., March, J.S., Schulte, A., Osborne, S., et al. (1994). Multimodality therapies: Methodologic issues in research and practice. *Child and Adolescent Psychiatry Clinics of North America, 3*, 361–377.
58. Wolraich, M.L. (2002) Current assessment and treatment practices in ADHD. In P.S. Jensen & J.R. Cooper (Eds.), *Attention deficit hyperactivity disorder: State of the science, best practices* (pp. 23–1–12). Kingston, NJ: Civic Research Institute.
59. Chronis, A.M., Fabiano, G.A., Gnagy, E.M., Onyango, A.N., Pelham, W.E., Williams, A., et al. (in press). An evaluation of the summer treatment program for children with attention-deficit/hyperactivity disorder using a treatment withdrawal design. *Behavior Therapy*.
60. Pelham, W. E. & Hoza, B. (1996). Intensive treatment: A summer treatment program for children with AD/HD. In E. Hibbs & P. Jensen (Eds.), *Psychosocial treatments for child and adolescent disorders: Empirically based strategies for clinical practice*. (pp. 311–340). New York: APA Press.
61. Pelham W.E., Greiner, A.R., & Gnagy, E.M. (1997). *Children's summer treatment program manual*. Buffalo, NY: Comprehensive Treatment for Attention Deficit Disorder.
62. Hoza, B., Mrug, S., Pelham, W.E., Jr., Greiner, A.R., & Gnagy, E.M. A friendship intervention for children with attention-deficit/hyperactivity disorder: Preliminary findings. *Journal of Attention Disorders, 6*, 87–98.
63. Mrug, S., Hoza, B., Gerdes, A. C. (2001). Children with attention-deficit/ hyperactivity disorder: Peer relationships and peer-oriented interventions. In D.W. Nangle & C.A. Erdley (Eds.), *The role of friendship in psychological adjustment: New directions for child and adolescent development* (pp. 51–77). San Francisco: Jossey-Bass.
64. MTA Cooperative Group. (1999). A 14-Month randomized clinical trial of treatment strategies for attention-deficit/hyperactivity disorder. *Archives of General Psychiatry, 56*, 1073–1086.
65. MTA Cooperative Group. (1999). Moderators and mediators of treatment response for children with attention-deficit/hyperactivity disorder. *Archives of General Psychiatry, 56*, 1088–1096.
66. Swanson, J.M., Kraemer, H.C., Hinshaw, S.P., Arnold, L.E., Conners, C.K., Abikoff, H.B., et al. Clinical relevance of the primary findings of the MTA: Success rates based on severity of ADHD and ODD symptoms at the end of treatment. *Journal of the American Academy of Child and Adolescent Psychiatry, 40*, 168–179.
67. Atkins, M.S., Pelham, W.E., & White, K.J. (1989). Hyperactivity and attention deficit disorder. In M. Hersen (Ed.), *Psychological aspects of developmental and physical disabilities: A casebook* (pp. 137–156). Thousand Oaks, CA: Sage.
68. Carlson, C.L., Pelham, W.E., Milich, R., & Dixon, J. (1992). Single and combined effects of methylphenidate and behavior therapy on the classroom performance of children with attention-deficit hyperactivity disorder. *Journal of Abnormal Child Psychology, 20*, 213–232.
69. Hinshaw, S.P., Heller, T., & McHale, J.P. (1992). Covert antisocial behavior in boys with attention deficit hyperactivity disorder: External validation and effects of methylphenidate. *Journal of Consulting and Clinical Psychology, 60*, 274–281.
70. Pelham, W.E., Schnedler, R.W., Bologna, N., & Contreras, A. (1980). Behavioral and stimulant treatment of hyperactive children: A therapy study with methylphenidate probes in a within-subject design. *Journal of Applied Behavioral Analysis, 13*, 221–236.
71. Pelham, W.E., Schnedler, R.W., Bender, M.E., Miller, J., Nilsson, D., Budrow, M., et al. (1988). The combination of behavior therapy and methylphenidate in the treatment of hyperactivity: A therapy outcome study. In L. Bloomingdale (Ed.), *Attention deficit disorder* (Vol. 3, pp. 29–48). London: Pergamon Press.
72. Barkley, R.A., & Murphy, K.R. (1998). *Attention-deficit hyperactivity disorder: A clinical workbook*. (2nd ed.). New York: Guilford.
73. Kendall, P.C., Flannery-Schroeder, E., Panichelli-Mindel, S.M., Southam-Gerow, M., Henin, A., & Warman, M. (1997). Therapy for youths with anxiety disorders: A second randomized clinical trial. *Journal of Consulting and Clinical Psychology, 65*(3), 366–380.
74. Clarke, G.N., Rhode, P., Lewinsohn, P.M., Hops, H., & Seeley, J.R. (1999). Cognitive-behavioral treatment of adolescent depression: Efficacy of acute group treatment and booster sessions. *Journal of the American Academy of Child and Adolescent Psychiatry, 38*, 272–279.

The information provided in this sheet was supported by Grant/Cooperative Agreement Number 1U84DD001049-01 from the Centers for Disease Control and Prevention (CDC). The contents are solely the responsibility of the authors and do not necessarily represent the official views of CDC. This fact sheet was approved by CHADD's Professional Advisory Board in 2004.

This fact sheet was updated in February 2004.

© 2004 Children and Adults with Attention-Deficit/Hyperactivity Disorder (CHADD).

For further information about AD/HD or CHADD, please contact:

**National Resource Center on AD/HD
Children and Adults with
Attention-Deficit/Hyperactivity Disorder**
4601 Presidents Drive, Suite 300
Lanham, MD 20706
800-233-4050
www.help4adhd.org

Please also visit the CHADD Web site at
www.chadd.org.

Appendix A

Classroom Behavioral Management Procedures

The following list includes typical classroom behavioral management procedures. They are arranged in order from mildest and least restrictive to more intensive and most restrictive procedures. Some of these programs may be included in 504 plans or Individualized Educational Programs for children with AD/HD (see CHADD Fact Sheet #4). Typically, an intervention is individualized and consists of several components based on the child's needs, classroom resources, and the teacher's skills and preferences.

1. Classroom rules and structure

- Use classroom rules such as:
 - Be respectful of others.
 - Obey adults.
 - Work quietly.
 - Stay in assigned seat/area.
 - Use materials appropriately.
 - Raise hand to speak or ask for help.
 - Stay on task and complete assignments.
- Post the rules and review them before each class until learned.
- Make rules objective and measurable.
- Tailor the number of rules to developmental level.
- Establish a predictable environment.
- Enhance children's organization (folders/charts for work).
- Evaluate rule-following and give feedback/consequences consistently.
- Tailor the frequency of feedback to developmental level.

2. Praise of appropriate behaviors and choosing battles carefully

- Ignore mild inappropriate behaviors that are not reinforced by peer attention.
- Use at least five times as many praises as negative comments.
- Use commands/reprimands to cue positive comments for children who are behaving appropriately — that is, find children who can be praised each time a reprimand or command is given to a child who is misbehaving.

3. Appropriate commands and reprimands

- Use clear, specific commands.
- Give private reprimands at the child's desk as much as possible.
- Reprimands should be brief, clear, neutral in tone, and as immediate as possible.

4. Individual accommodations and structure for the child

- Structure the classroom to maximize the child's success.
- Place the student's desk near the teacher to facilitate monitoring.
- Enlist a peer to help the student copy assignments from the board.
- Break assignments into small chunks.
- Give frequent and immediate feedback.
- Require corrections before new work is given.

5. Proactive interventions to increase academic performance

- Such interventions can prevent problematic behavior from occurring and can be implemented by individuals other than the classroom teacher, such as peers or a classroom aide. When disruptive behavior is not the primary problem, these academic interventions can improve behavior significantly.
- Focus on increasing completion and accuracy of work.
 - Offer task choices.
 - Provide peer tutoring.
 - Consider computer-assisted instruction.

6. "When...then" contingencies (withdrawing rewards or privileges in response to inappropriate behavior)—Examples include recess time contingent upon completion of work, staying after school to complete work, assigning less desirable work prior to more desirable assignments, and requiring assignment completion in study hall before allowing free time.

7. Daily school-home report card (instruction packet available at <http://wings.buffalo.edu/adhd>)—This tool allows parents and teacher to communicate regularly, identifying, monitoring and changing classroom

problems. It is inexpensive and minimal teacher time is required.

- Teachers determine the individualized target behaviors.
- Teachers evaluate targets at school and send the report card home with the child.
- Parents provide home-based rewards; more rewards for better performance and fewer for lesser performance.
- Teachers continually monitor and make adjustments to targets and criteria as behavior improves or new problems develop.
- Use the report card with other behavioral components such as commands, praise, rules, and academic programs.

8. Behavior chart and/or reward and consequence program (point or token system)

- Establish target behaviors and ensure that the child knows the behaviors and goals (e.g., list on index card taped to desk).
- Establish rewards for exhibiting target behaviors.
- Monitor the child and give feedback.
- Reward young children immediately.
- Use points, tokens or stars that can later be exchanged for rewards.

9. Classwide interventions and group contingencies

—Such interventions encourage children to help one another because everyone can be rewarded. There is also potential for improvement in the behavior of the entire class.

- Establish goals for the class as well as the individual.
- Establish rewards for appropriate behavior that any student can earn (e.g., class lottery, jelly bean jar, wacky bucks).
- Establish a class reward system in which the entire class (or subset of the class) earns rewards based on class functioning as a whole (e.g., Good Behavior Game) or the functioning of the student with AD/HD.
- Tailor frequency of rewards and consequences to developmental level.

10. Time out—The child is removed, either in the classroom or to the office, from the ongoing activity for a few minutes (less for younger children and more for older) when he or she misbehaves.

11. Schoolwide programs—Such programs, which include schoolwide discipline plans, can be structured to minimize the problems experienced by children with AD/HD, while at the same time help manage the behavior of all students in a school.

Sample Daily Report Card

Child's Name: _____ Date: _____

	Special		Language Arts		Math	Reading	Science	
	Y	N	Y	N			Y	N
Follows class rules (no more than 3 rule violations per period).	Y	N	Y	N	Y	N	Y	N
Completes assignments within the designated time.	Y	N	Y	N	Y	N	Y	N
Completes assignments with 80% accuracy.	Y	N	Y	N	Y	N	Y	N
Complies with teacher requests (no more than 3 instances of noncompliance per period).	Y	N	Y	N	Y	N	Y	N
No more than 3 instances of teasing per period.	Y	N	Y	N	Y	N	Y	N
Follows lunch rules (no more than 3 violations).	Y	N						
Follows recess rules (no more than 3 violations).	Y	N						
Other	Y	N						

Total Number of "Yes" Answers _____

Teacher's Initials: _____

Comments: _____